Vyalev (Foslevodopa/Foscarbidopa) and Vyafuser (AbbVie Infusion Pump)

The patient and support person know this medication and the pump best

The medication and infusion pump are used for symptom management of Parkinson's disease. The pump delivers Vyalev (Foslevodopa/Foscarbidopa) subcutaneously 24hrs a day.

If this is a medical emergency, stabilize the patient before addressing the pump

Vyalev (Foslevodopa / Foscarbidopa) should remain infusing, except in the following situations:

- **Suspected cellulitis related to the medication infusion**
- **Patient is no longer able to maintain the infusion (ie. due to loss of consciousness)**
- **MRI (the pump is not MRI compatible, while the infusion set is. The infusion must be stopped and disconnected prior to the scan)**
- **Psychosis**

Once the pump is stopped, oral medication must be started within ONE (1) hour

If infusion has been stopped for more than ONE (1) hour, infusion set must be replaced

If the patient or support person is not able, STOP the pump using the following instructions:

1. On the main STATUS SCREEN, press the MENU BUTTON

2. In the MENU SCREEN, use the UP ARROW (a) to highlight STOP PUMP.

3. Once STOP PUMP is highlighted, press SELECT, then YES

For additional instructions for management of Vyalev or Vyafuser, including (re)initiating oral medication, please contact:

Movement Disorder Clinic **Deer Lodge Centre**	**204-940-8400** **Monday to Friday 8am-4pm**
Movement Disorder Clinic **Clinical Resource Nurse** **Kelly Williams**	**204-940-8427** **kwilliams3@mdc-dlc.ca**
Rechelle Flores **AbbVie Home Nurse**	**1-844-686-3672 ext. 5122** **431-323-2324**

MOVEMENT DISORDER CLINIC
DEER LODGE CENTRE
Making lives better

200 Woodlawn Street, Winnipeg, Manitoba, R3J 2H7
Phone: 204-940-8400 Fax: 204-940-8414